

Supplementary Table. Source of life history parameters for the reef fish species included in the meta-analysis.

Family	Scientific name	Longevity source	Growth source	Maturity source
Haemulidae	<i>Diagramma pictum</i>		(Baillon, 1994)	(Grandcourt et al., 2011)
Haemulidae	<i>Haemulon album</i>		(Gaut and Munro, 1983)	(Gaut and Munro, 1983)
Haemulidae	<i>Haemulon parra</i>		(Valle et al., 1997)	
Haemulidae	<i>Haemulon plumieri</i>	(Araújo and Martins, 2007)	(Araújo and Martins, 2007)	(Araújo and Martins, 2007)
Haemulidae	<i>Haemulon sciurus</i>	(Pitt et al., 2010)	(Pitt et al., 2010)	
Haemulidae	<i>Microlepidotus brevipinnis</i>	(Espino-Barr et al., 2010)	(Espino-Barr et al., 2010)	
Haemulidae	<i>Orthopristis ruber</i>	(Guerrieri et al., 2015)	(Guerrieri et al., 2015)	
Haemulidae	<i>Parapristipoma trilineatum</i>	(Doiuchi et al., 2007)	(Doiuchi et al., 2007)	
Haemulidae	<i>Plectorhinchus pictus</i>	(Wenjia et al., 2015)	(Wenjia et al., 2015)	
Haemulidae	<i>Plectorhinchus sordidus</i>	(Grandcourt et al., 2011)	(Grandcourt et al., 2011)	(Grandcourt et al., 2011)
Haemulidae	<i>Pomadasys argenteus</i>	(Mathews and Samuel, 1991)	(Kulbicki et al., 2009)	
Haemulidae	<i>Pomadasys commersonnii</i>		(Torres Jr. and Pauly, 1991)	
Haemulidae	<i>Pomadasys kaakan</i>	(Al-Husaini et al., 2002)	(Al-Husaini et al., 2002)	(Al-Husaini et al., 2002)
Haemulidae	<i>Pomadasys maculatus</i>	(Bade, 1989)	(Bade, 1989)	(Bade, 1989)
Labridae	<i>Bodianus axillaris</i>	(Hubble, 2003)	(Hubble, 2003)	
Labridae	<i>Bodianus frenchii</i>	(Cossington et al., 2010)	(Cossington et al., 2010)	(Cossington et al., 2010)
Labridae	<i>Bodianus mesothorax</i>	(Hubble, 2003)	(Hubble, 2003)	
Labridae	<i>Choerodon cauteroma</i>	(Fairclough, 2005)	(Fairclough, 2005)	(Fairclough, 2005)
Labridae	<i>Choerodon cyanodus</i>	(Fairclough, 2005)	(Fairclough, 2005)	(Fairclough, 2005)
Labridae	<i>Choerodon fasciatus</i>	(Hubble, 2003)	(Hubble, 2003)	
Labridae	<i>Choerodon rubescens</i>	(Nardi et al., 2006)	(Fairclough, 2005)	(Fairclough, 2005)
Labridae	<i>Choerodon schoenleinii</i>	(Ebisawa et al., 2010)	(Ebisawa et al., 2010)	(Ebisawa et al., 2010)
Labridae	<i>Cheilinus undulatus</i>	(Choat et al., 2006)	(Sadovy et al., 2007)	(Sadovy et al., 2007)
Labridae	<i>Cheilinus fasciatus</i>	(Hubble, 2003)	(Hubble, 2003)	
Labridae	<i>Lachnolaimus maximus</i>	(McBride and Richardson, 2007)	(McBride and Richardson, 2007)	(McBride et al., 2008)
Labridae	<i>Thalassoma lunare</i>	(Ackerman, 2004)	(Ackerman, 2004)	(Ackerman, 2004)
Shark-Lamnidae	<i>Isurus oxyrinchus</i>	(Semba et al., 2009)	(Semba et al., 2009)	(Semba et al., 2009)

Family	Scientific name	Longevity source	Growth source	Maturity source
Serranidae	<i>Aethaloperca rogaa</i>	(Mapleston et al., 2009)	(Mapleston et al., 2009)	
Serranidae	<i>Anyperodon leucogrammicus</i>	(Mapleston et al., 2009)	(Mapleston et al., 2009)	
Serranidae	<i>Cephalopholis argus</i>	(Donovan et al., 2013)	(Donovan et al., 2013)	
Serranidae	<i>Cephalopholis boenak</i>	(Chan and Sadovy, 2002)	(Chan and Sadovy, 2002)	(Chan and Sadovy, 2002)
Serranidae	<i>Cephalopholis cruentata</i>	(Potts and Manooch, 1999)	(Potts and Manooch, 1999)	
Serranidae	<i>Cephalopholis fulva</i>	(Trott, 2006)	(Trott, 2006)	
Serranidae	<i>Cephalopholis miniata</i>	(Mathews and Samuel, 1991)	(Mathews and Samuel, 1991)	
Serranidae	<i>Cephalopholis panamensis</i>	(Craig et al., 1999)	(Craig et al., 1999)	(Erisman et al., 2010)
Serranidae	<i>Cephalopholis spiloparaea</i>	(Fry et al., 2006)	(Fry et al., 2006)	
Serranidae	<i>Cephalopholis taeniops</i>	(Tariche et al., 2015)	(Tariche et al., 2015)	(Pastor et al., 2002)
Serranidae	<i>Cephalopholis urodetta</i>	(Fry et al., 2006)	(Fry et al., 2006)	
Serranidae	<i>Cromileptes altivelis</i>	(Williams et al., 2009)	(Williams et al., 2009)	
Serranidae	<i>Diplectrum formosum</i>	(Bubley and Pashuk, 2010)	(Bubley and Pashuk, 2010)	(Bubley and Pashuk, 2010)
Serranidae	<i>Diplectrum labarum</i>		(Morales-Nin, 1994)	(Foster and Vincent, 2010)
Serranidae	<i>Diplectrum macropoma</i>		(Morales-Nin, 1994)	
Serranidae	<i>Epinephelus adscensionis</i>	(Burton et al., 2012)	(Burton et al., 2012)	
Serranidae	<i>Epinephelus areolatus</i>	(Mathews and Samuel, 1991)	(Mathews and Samuel, 1991)	(Mahmoud and others, 2009)
Serranidae	<i>Epinephelus chlorostigma</i>	(Grandcourt, 2002)	(Grandcourt, 2002)	(Mees, 1992)
Serranidae	<i>Epinephelus coioides</i>	(Grandcourt et al., 2005)	(Grandcourt et al., 2005)	(Grandcourt et al., 2005)
Serranidae	<i>Epinephelus cyanopodus</i>	(Loubens, 1980)	(Loubens, 1980)	
Serranidae	<i>Epinephelus diacanthus</i>	(Mehanna et al., 2013)	(Edwards and Shaher, 1991)	
Serranidae	<i>Epinephelus fuscoguttatus</i>	(Grandcourt, 2005)	(Grandcourt, 2005)	(Pears et al., 2006)
Serranidae	<i>Epinephelus guttatus</i>	(Sadovy et al., 1992)	(Sadovy et al., 1992)	(Sadovy et al., 1992)
Serranidae	<i>Epinephelus itajara</i>	(Bullock et al., 1992)	(Bullock et al., 1992)	
Serranidae	<i>Epinephelus labriformis</i>	(Craig et al., 1999)	(Craig et al., 1999)	
Serranidae	<i>Epinephelus latifasciatus</i>	(Mathews and Samuel, 1991)	(Mathews and Samuel, 1991)	(Sujatha and Shrikanya, 2013)
Serranidae	<i>Epinephelus maculatus</i>	(Loubens, 1980)	(Loubens, 1980)	(Loubens, 1980)

Family	Scientific name	Longevity source	Growth source	Maturity source
Serranidae	<i>Epinephelus merra</i>	(Loubens, 1980)	(Pothin et al., 2004)	
Serranidae	<i>Epinephelus morio</i>	(Lombardi-Carlson et al., 2002)	(Lombardi-Carlson et al., 2002)	(Collins et al., 2002)
Serranidae	<i>Epinephelus morrhua</i>	(Fry et al., 2006)	(Fry et al., 2006)	
Serranidae	<i>Epinephelus multinotatus</i>	(Grandcourt, 2005)	(Grandcourt, 2005)	
Serranidae	<i>Epinephelus ongus</i>	(Craig, 2007)	(Ohta and Ebisawa, 2016)	
Serranidae	<i>Epinephelus polyphekadion</i>	(Grandcourt, 2005)	(Rhodes et al., 2011)	(Rhodes et al., 2011)
Serranidae	<i>Epinephelus quoyanus</i>	(Mapleston et al., 2009)	(Mapleston et al., 2009)	(Al-Salamah, 2014)
Serranidae	<i>Epinephelus radiatus</i>	(Mahé et al., 2016)	(Mahé et al., 2016)	
Serranidae	<i>Epinephelus rivulatus</i>	(Loubens, 1980)	(Loubens, 1980)	(Mackie, 2000)
Serranidae	<i>Epinephelus striatus</i>	(Bush et al., 2006)	(Bush et al., 2006)	(Sadovy and Colin, 1995)
Serranidae	<i>Epinephelus tauvina</i>	(Mathews and Samuel, 1991)	(Mathews and Samuel, 1991)	(El-Sayed and Abdel-Bary, 1999)
Serranidae	<i>Epinephelus tukula</i>	(Grandcourt, 2005)	(Grandcourt, 2005)	
Serranidae	<i>Hyporthodus mystacinus</i>	(Farmer et al., 2016)	(Farmer et al., 2016)	(Farmer et al., 2016)
Serranidae	<i>Hyporthodus nigritus</i>	(Manooch, 1987)	(Manooch, 1987)	(Farmer et al., 2016)
Serranidae	<i>Hyporthodus niveatus</i>	(Kowal, 2010)	(Kowal, 2010)	(Wyanski et al., 2000)
Serranidae	<i>Hyporthodus octofasciatus</i>	(Wakefield et al., 2013)	(Wakefield et al., 2013)	(Wakefield et al., 2013)
Serranidae	<i>Hyporthodus quernus</i>	(Nichols and DeMartini, 2008)	(Nichols and DeMartini, 2008)	(DeMartini et al., 2011)
Serranidae	<i>Mycteroperca bonaci</i>	(Crabtree and Bullock, 1998)	(Crabtree and Bullock, 1998)	(Crabtree and Bullock, 1998)
Serranidae	<i>Mycteroperca fusca</i>	(Bustos et al., 2009)	(Bustos et al., 2009)	(Bustos et al., 2010)
Serranidae	<i>Mycteroperca interstitialis</i>	(Manickchand-Heileman and Phillip, 2000)	(Burton et al., 2014)	
Serranidae	<i>Mycteroperca microlepis</i>	(Hood and Schlieder, 1992)	(Hood and Schlieder, 1992)	(McGovern et al., 1998)
Serranidae	<i>Mycteroperca olfax</i>	(Usseglio et al., 2015)	(Usseglio et al., 2015)	(Usseglio et al., 2015)
Serranidae	<i>Mycteroperca phenax</i>	(Harris et al., 2002)	(Harris et al., 2002)	(Harris et al., 2002)
Serranidae	<i>Mycteroperca venenosa</i>	(Burton et al., 2015)	(Burton et al., 2015)	
Serranidae	<i>Paralabrax maculatofasciatus</i>	(Andrews et al., 2005)	(Andrews et al., 2005)	(Allen et al., 1995)

Family	Scientific name	Longevity source	Growth source	Maturity source
Serranidae	<i>Plectropomus laevis</i>	(Grandcourt, 2005)	(Heupel et al., 2010)	(Heupel et al., 2010)
Serranidae	<i>Plectropomus leopardus</i>	(Loubens, 1980)	(Ferreira, 1995)	(Ferreira, 1995)
Serranidae	<i>Plectropomus maculatus</i>	(Ferreira and Russ, 1992)	(Ferreira and Russ, 1992)	(Ferreira, 1993)
Serranidae	<i>Pronotogrammus martinicensis</i>	(McBride et al., 2009)	(McBride et al., 2009)	
Serranidae	<i>Variola albimarginata</i>	(Mapleston et al., 2009)	(Mapleston et al., 2009)	
Serranidae	<i>Variola louti</i>	(Grandcourt, 2005)	(Grandcourt, 2005)	
Shark-Alopiidae	<i>Alopias pelagicus</i>	(Drew et al., 2015)	(Drew et al., 2015)	(Drew et al., 2015)
Shark-Alopiidae	<i>Alopias superciliosus</i>	(Fernandez-Carvalho et al., 2011)	(Liu et al., 1998)	
Shark-Carcharhinidae	<i>Carcharhinus amblyrhynchos</i>	(Robbins, 2006)	(Robbins, 2006)	(Robbins, 2006)
Shark-Carcharhinidae	<i>Carcharhinus cautus</i>	(White et al., 2002)	(White et al., 2002)	(White et al., 2002)
Shark-Carcharhinidae	<i>Carcharhinus falciformis</i>	(Bonfil et al., 1993)	(Bonfil et al., 1993)	(Branstetter, 1987)
Shark-Carcharhinidae	<i>Carcharhinus leucas</i>	(Neer et al., 2005)	(Neer et al., 2005)	
Shark-Carcharhinidae	<i>Carcharhinus limbatus</i>	(Killam and Parsons, 1989)	(Killam and Parsons, 1989)	(Castro, 1996)
Shark-Carcharhinidae	<i>Carcharhinus melanopterus</i>	(Chin et al., 2013)	(Chin et al., 2013)	(Chin et al., 2013)
Shark-Carcharhinidae	<i>Carcharhinus obscurus</i>	(Simpfendorfer et al., 2002)	(Simpfendorfer et al., 2002)	(Simpfendorfer et al., 2002)
Shark-Carcharhinidae	<i>Carcharhinus plumbeus</i>	(Romine et al., 2006)	(Romine et al., 2006)	(Romine et al., 2006)
Shark-Carcharhinidae	<i>Carcharhinus sorrah</i>	(Davenport and Stevens, 1988)	(Davenport and Stevens, 1988)	(Harry et al., 2013)
Shark-Carcharhinidae	<i>Carcharhinus tilstoni</i>	(Davenport and Stevens, 1988)	(Davenport and Stevens, 1988)	(Harry et al., 2013)
Shark-Carcharhinidae	<i>Loxodon macrorhinus</i>	(Gutteridge et al., 2013)	(Gutteridge et al., 2013)	(Gutteridge et al., 2013)

Family	Scientific name	Longevity source	Growth source	Maturity source
Shark-Carcharhinidae	<i>Negaprion brevirostris</i>	(Brown and Gruber, 1988)	(Brown and Gruber, 1988)	(Gruber and Pulos, 1982)
Shark-Carcharhinidae	<i>Rhizoprionodon acutus</i>	(Ba et al., 2015)	(Ba et al., 2015)	(Henderson et al., 2006)
Shark-Carcharhinidae	<i>Rhizoprionodon porosus</i>	(Lessa and Santana, 1998)	(Lessa and Santana, 1998)	(Mattos et al., 2001)
Shark-Carcharhinidae	<i>Rhizoprionodon taylori</i>	(Simpfendorfer, 1993)	(Simpfendorfer, 1993)	(Simpfendorfer, 1992)
Shark-Carcharhinidae	<i>Rhizoprionodon terraenovae</i>	(Loefer and Sedberry, 2003)	(Parsons, 1985)	
Shark-Carcharhinidae	<i>Triaenodon obesus</i>	(Robbins, 2006)	(Robbins, 2006)	(Robbins, 2006)
Shark-Sphyrnidae	<i>Sphyraña lewini</i>	(Piercey et al., 2007)	(Piercey et al., 2007)	(Liu and Chen, 1999)
Shark-Sphyrnidae	<i>Sphyraña mokarran</i>	(Piercey et al., 2010)	(Piercey et al., 2010)	(Piercey et al., 2010)
Shark-Sphyrnidae	<i>Sphyraña tiburo</i>	(Carlson and Parsons, 1997)	(Parsons, 1993)	(Lombardi-Carlson et al., 2003)

## References

- Ackerman, J.  
2004. Geographic variation in size at age of the coral reef fish, *Thalassoma Lunare* (Family: Labridae): A contribution to life history theory. Ph.D diss. James Cook University, Australia.
- Al-Husaini, M., Al-Baz, A., Al-Ayoub, S., Safar, S., Al-Wazan, Z., and S., Al-Jazzaf  
2002. Age, growth, mortality, and yield-per-recruit for nagroor, *Pomadasys kakaan*, in Kuwait's waters. Fish. Res. 59:101–115
- Allen, L. G., Hovey, T. E., Love, M. S., and J. T., Smith  
1995. The life history of the spotted sand bass (*Paralabrax maculatofasciatus*) within the southern California bight. Calif. Coop. Ocean. Fish. Investig. Rep. 193–203
- Al-Salamah, M.  
2014. Connectivity of the longfin grouper (*Epinephelus Quoyanus*) in a marine reserve in the Great Keppel Island Group. M.S. thesis. King Abdullah University of science and Technology.
- Andrews, A. H., Laman, E. A., Bennett, T., Jones, E., and G. M., Cailliet.  
2005. Age and growth of spotted sand bass, *Paralabrax maculatofasciatus*, in Bahia de Los Angeles, Baja California, Mexico, with age validation using otolith edge analysis. Bull. South. Calif. Acad. Sci. 104:14–25
- Araújo, J. N., and A. S., Martins.  
2007. Age, growth and mortality of white grunt (*Haemulon plumieri*) from the central coast of Brazil. Sci. Mar. 71:793–800
- Ba, A., Diouf, K., Guilhaumon, F., and J., Panfili.  
2015. Slow growth of the overexploited milk shark *Rhizoprionodon acutus* affects its sustainability in West Africa. J. Fish Biol. 87:912–929. <https://doi.org/10.1111/jfb.12764>.
- Bade, T. M.  
1989. Aspects of the biology of grunts (Teleostei : Haemulidae) from North Queensland waters. Ph.D diss. James Cook University, Australia.

Baillon, N.  
1994. Lecture des stries annuelles sur les otolithes de *Diagramma pictum*. Pages 211–238  
Otolithometrie en milieu tropical: Application a trois especes Du Lagon De Nouvelle-  
Caledonie, 1st edition. Institut Francais de Recherche Scientifique Pour le Developpement en  
Cooperation, Paris.

Bonfil, R., Mena, R., and D., de Anda.  
1993. Biological Parameters of Commercially Exploited silky sharks, *Carcharhinus falciformis*, from the Campeche Bank, Mexico (Technical Report). Campeche Bank, Mexico:  
NOAA.

Branstetter, S.  
1987. Age, growth and reproductive biology of the silky shark, *Carcharhinus falciformis*, and  
the scalloped hammerhead, *Sphyrna lewini*, from the northwestern Gulf of Mexico. Environ.  
Biol. Fishes 19:161–173

Brown, C. A., and S. H., Gruber.  
1988. Age assessment of the lemon shark, *Negaprion brevirostris*, using tetracycline validated  
vertebral Centra. Copeia 1988:747. <https://doi.org/10.2307/1445397>.

Bubley, W. J., and O., Pashuk.  
2010. Life history of a simultaneously hermaphroditic fish, *Diplectrum formosum*. J. Fish  
Biol. 77:676-691. <https://doi.org/10.1111/j.1095-8649.2010.02710.x>.

Bullock, L. H., Murphy, M. D., Godcharles, M. F., and M. E. Mitchell.  
1992. Age, growth, and reproduction of jewfish *Epinephelus itajara* in the eastern Gulf of  
Mexico. Fish. Bull.

Burton, M. L., Potts, J. C., and D. R., Carr.  
2012. Age, growth, and natural mortality of rock hind, *Epinephelus adscensionis*, from the  
Gulf of Mexico. Bull. Mar. Sci. 88:903–917. <https://doi.org/10.5343/bms.2011.1102>.

Burton, M. L., Potts, J. C., and D. R., Carr.  
2014. Age, growth, and mortality of yellowmouth grouper from the Southeastern United  
States. Mar. Coast. Fish. 6:33–42. <https://doi.org/10.1080/19425120.2013.866998>.

Burton, M. L., Potts, J. C., and D. R., Carr.  
2015. Age, growth, and natural mortality of yellowfin grouper (*Mycteroperca venenosa*) from the southeastern United States. PeerJ 3:e1099. <https://doi.org/10.7717/peerj.1099>.

Bush, P. G., Lane, E. D., Ebanks-Petrie, G. C., Luke, K., Johnson, B., McCoy, C., Bothwell, J., and E., Parsons.  
2006. The Nassau grouper spawning aggregation fishery of the Cayman Islands—an historical and management perspective. 57th Gulf and Caribbean Fisheries Institute. 515-524.

Bustos, R., Luque, Á., and J. G., Pajuelo.  
2009. Age estimation and growth pattern of the island grouper, *Mycteroperca fusca* (Serranidae) in an island population on the northwest coast of Africa. Sci. Mar. 73:319–328. <https://doi.org/10.3989/scimar.2009.73n2319>.

Bustos, R., Luque, Á., and J. G., Pajuelo.  
2010. Reproductive biology of the island grouper (*Mycteroperca fusca*) in the Canary Islands, northwest coast of Africa. Sci. Mar. 74:613–619. <https://doi.org/10.3989/scimar.2010.74n3613>.

Carlson, J. K., and G. R., Parsons.  
1997. Age and growth of the bonnethead shark, *Sphyrna tiburo*, from northwest Florida, with comments on clinal variation. Environ. Biol. Fishes 50:331–341

Castro, J. I.  
1996. Biology of the blacktip shark, *Carcharhinus limbatus*, off the southeastern United States. Bull. Mar. Sci. 59:508–522

Chan, T. T., and Y. Sadovy.  
2002. Reproductive biology, age and growth in the chocolate hind, *Cephalopholis boenak* (Bloch, 1790), in Hong Kong. Mar. Freshw. Res. 53:791–803

Chin, A., Simpfendorfer, C., Tobin, A., and M., Heupel.  
2013. Validated age, growth and reproductive biology of *Carcharhinus melanopterus*, a widely distributed and exploited reef shark. Mar. Freshw. Res. 64:965. <https://doi.org/10.1071/MF13017>.

Choat, J. H., Davies, C. R., Ackerman, J. L., and B. D. Mapstone.  
2006. Age structure and growth in a large teleost, *Cheilinus undulatus*, with a review of size distribution in labrid fishes. Mar. Ecol. Prog. Ser. 318:237–246

Collins, L. A., Fitzhugh, G. R., Lombardi-Carlson, L. A., Lyon, H. M., Walling, W. T., and D. W., Oliver.  
2002. Characterization of red grouper (Serranidae: *Epinephelus morio*) reproduction from the eastern Gulf of Mexico. Natl. Mar. Fish. Serv. Panama City Lab. Contrib. Ser. 7

Cossington, S., Hesp, S. A., Hall, N. G., and I. C., Potter.  
2010. Growth and reproductive biology of the foxfish *Bodianus frenchii*, a very long-lived and monandric protogynous hermaphroditic labrid. J. Fish Biol. 77(3):600–626.  
<https://doi.org/10.1111/j.1095-8649.2010.02706.x>.

Crabtree, R. E., and L. H., Bullock.  
1998. Age, growth, and reproduction of black grouper, *Mycteroperca bonaci*. Fla. Waters Fish Bull 96:735–753

Craig, M. T.  
2007. Preliminary observations on the life history of the white-streaked grouper, *Epinephelus ongus*, from Okinawa, Japan. Ichthyol. Res. 54:81–84. <https://doi.org/10.1007/s10228-006-0377-8>.

Craig, M. T., Pondella, D. J., and J. C., Hafner.  
1999. Analysis of age and growth in two eastern pacific groupers (Serranidae: Epinephelinae). Bull. Mar. Sci. 65:807–814

Davenport, S., and J. D., Stevens.  
1988. Age and growth of two commercially imported sharks (*Carcharhinus tilstoni* and *C. sorrah*) from Northern Australia. Mar. Freshw. Res. 39:417–433

DeMartini, E. E., Everson, A. R., and R.S., Nichols.  
2011. Estimates of body sizes at maturation and at sex change, and the spawning seasonality and sex ratio of the endemic Hawaiian grouper (*Hoporthodus quernus*, F. Epinephelidae). Fish. Bull. 109:123–134

Doiuchi, R., Kokubo, T., and M. Ogawa.  
2007. Age and growth of threeline grunt *Parapristipoma trilineatum* along the south-western coast of Kii Peninsula, Japan. Fish. Sci. 73:777–783. <https://doi.org/10.1111/j.1444-2906.2007.01396.x>.

Donovan, M. K., Friedlander, A. M., DeMartini, E. E., Donahue, M. J., and I.D. Williams.  
2013. Demographic patterns in the peacock grouper (*Cephalopholis argus*), an introduced Hawaiian reef fish. Environ. Biol. Fishes 96:981–994. <https://doi.org/10.1007/s10641-012-0095-1>.

Drew, M., White, W. T., Dharmadi, D., Harry, A.V., and C., Huvemeers.  
2015. Age, growth and maturity of the pelagic thresher *Alopias pelagicus* and the scalloped hammerhead *Sphyrna lewini*: age and growth of two large shark species. J. Fish Biol. 86:333–354. <https://doi.org/10.1111/jfb.12586>.

Ebisawa, A., Kanashiro, K., and T., Kiyan.  
2010. Growth, sex ratio, and maturation rate with age in the blackspot tuskfish *Choerodon schoenleinii* in waters off Okinawa Island, southwestern Japan. Fish. Sci. 76:577–583. <https://doi.org/10.1007/s12562-010-0244-4>.

Edwards, R., and S. Shaher.  
1991. Biometrics of Marine Fishes from the Gulf of Aden. Fishbyte

El-Sayed, A-F. M., and K., Abdel-Bary.  
1999. Reproductive biology of grouper fish *Epinephelus tauvina* (Family Serranidae) in the Arabian Gulf waters. Indian J. Mar. Sci. 28:89–91

Erisman, B. E., Craig, M. T., and P.A., Hastings.  
2010. Reproductive biology of the Panama graysby *Cephalopholis panamensis* (Teleostei: Epinephelidae). J. Fish Biol. 76:1312–1328. <https://doi.org/10.1111/j.1095-8649.2010.02567.x>.

Espino-Barr, E., Gallardo-Cabello, M., Granados-Flores, K., Cabral-Solis, E.G., Garcia-Boa, A., and M., Puente-Gomez.  
2010. Growth analysis of *Microlepidotus brevipinnis* from the Pacific Coast of Jalisco, Mexico. J. Fish. Aquat. Sci. 5:293–303

Fairclough, D.  
2005. The biology of four tuskfish species (Choerodon: Labridae) in Western Australia. Ph.D thesis. Murdoch University.

Farmer, N. A., Malinowski, R. P., McGovern, M. F., and P. J., Rubec.  
2016. Stock complexes for fisheries management in the Gulf of Mexico. Mar. Coast. Fish. 8:177–201. <https://doi.org/10.1080/19425120.2015.1024359>.

Fernandez-Carvalho, J., Coelho, R., Erzini, K., and M., Neves Santos.  
2011. Age and growth of the bigeye thresher shark, *Alopias superciliosus*, from the pelagic longline fisheries in the tropical northeastern Atlantic Ocean, determined by vertebral band counts. Aquat. Living Resour. 24:359–368. <https://doi.org/10.1051/alr/2011046>.

Ferreira, B. P.  
1993. Reproduction of the inshore coral trout *Plectropomus maculates* (Perciformes: Serranidae) from the central Great Barrier Reef, Australia. J. Fish Biol. 42:831–844

Ferreira, B. P.  
1995. Reproduction of the common coral trout *Plectropomus leopardus* (Serranidae: Epinephelinae) from the central and northern Great Barrier Reef, Australia. Bull. Mar. Sci. 56:653–669

Ferreira, B. P., and G.R., Russ.  
1992. Age, growth and mortality of the inshore coral trout *Plectropomus maculatus* (Pisces: Serranidae) from the Central Great Barrier Reef, Australia. Mar. Freshw. Res. 43:1301–1312

Foster, S. J., and A.C.J., Vincent.  
2010. Using life-history information to assess potential effects of shrimp trawling on small fishes. J. Fish Biol. 76:2434–2454. <https://doi.org/10.1111/j.1095-8649.2010.02631.x>.

Fry, G. C., Brewer, D. T., and W.N., Venables.  
2006. Vulnerability of deepwater demersal fishes to commercial fishing: Evidence from a study around a tropical volcanic seamount in Papua New Guinea. Fish. Res. 81:126–141. <https://doi.org/10.1016/j.fishres.2006.08.002>.

Gaut, V. C., and J. L., Munro.  
1983. The biology, ecology, and bionomics of the grunts, Pomadasysidae. In Caribb. Coral Reef Fish. Resour. p. 110–141. International Center for Living Aquatic Resources Management, Manila, Philippines., 2nd ed.

Grandcourt, E. M., Al Abdessalaam, T. Z., Francis, F., and A., Al Shamsi.  
2011. Demographic parameters and status assessments of *Lutjanus ehrenbergii*, *Lethrinus lentjan*, *Plectorrhinchus sordidus*, and *Rhabdosargus sarba* in the southern Arabian Gulf. J. Appl. Ichthyol. 27:1203–1211. <https://doi.org/10.1111/j.1439-0426.2011.01776.x>.

Grandcourt, E. M.  
2005. Demographic characteristics of selected epinepheline groupers (family Serranidae, subfamily Epinephelinae) from Aldabra Atoll, Seychelles. National Museum of Natural History, Smithsonian Institution Washington.

Grandcourt, E. M., Al Abdessalaam, T. Z., Francis, F., and A. T., Al Shamsi.  
2005. Population biology and assessment of the orange-spotted grouper, *Epinephelus coioides* (Hamilton, 1822), in the southern Arabian Gulf. Fish. Res. 74:55–68

Grandcourt, E. M., Al Abdessalaam, T. Z., Francis, F., and A. T., Al Shamsi.  
2011. Reproductive biology and implications for management of the painted sweetlips *Diagramma pictum* in the southern Arabian Gulf. J. Fish Biol. 79:615–632.  
<https://doi.org/10.1111/j.1095-8649.2011.03044.x>.

Grandcourt, E. M.  
2002. Demographic characteristics of a selection of exploited reef fish from the Seychelles: preliminary study. Mar. Freshw. Res. 53:123–130

Gruber, S. H., and L. M., Pulos.  
1982. Biological materials for the study of age and growth in a tropical marine Elasmobranch, the lemon shark, *Negaprion brevirostris* (poey). In Workshop Age Determ. Ocean. Pelagic Fishes Tunas Billfishes Sharks. p. 193–205. National Marine Fisheries Service, NOAA, Southeast Fisheries Science Center, Miami Laboratory.

Guerrieri, A., Eslava, N., González, L. W., and F., Guevara.  
2015. Growth and mortality parameters of *Orthopristis ruber* (Perciformes: Haemulidae) from Los Frailes Archipelago, Venezuela. Rev. Biol. Trop. 63:189–198

Gutteridge, A. N., Huveneers, C., Marshall, L. J., Tibbetts, I. R., and M. B., Bennett.  
2013. Life-history traits of a small-bodied coastal shark. Mar. Freshw. Res. 64:54.  
<https://doi.org/10.1071/MF12140>.

Harris, P. J., Wyanski, D. M., White, D. B., and J. L., Moore.  
2002. Age, growth, and reproduction of scamp, *Mycteroperca phenax*, in the southwestern North Atlantic, 1979–1997. Bull. Mar. Sci. 70:113–132

Harry, A. V., Tobin, A. J., and C. A., Simpfendorfer.  
2013. Age, growth and reproductive biology of the spot-tail shark, *Carcharhinus sorrah*, and the Australian blacktip shark, *C. tilstoni*, from the Great Barrier Reef World Heritage Area, north-eastern Australia. Mar. Freshw. Res. 64:277. <https://doi.org/10.1071/MF12142>.

Henderson, A. C., McIlwain, J. L., Al-Oufi, H. S., and A., Ambu-Ali.  
2006. Reproductive biology of the milk shark *Rhizoprionodon acutus* and the bigeye houndshark *Iago omanensis* in the coastal waters of Oman. J. Fish Biol. 68:1662–1678

Heupel, M. R., Williams, A. J., Welch, D. J., Davies, C. R., Adams, S., Carlos, G., and B. D., Mapstone.  
2010. Demography of a large exploited grouper, *Plectropomus laevis*: Implications for fisheries management. Mar. Freshw. Res. 61:184. <https://doi.org/10.1071/MF09056>.

Hood, P. B., and R. A., Schlieder.  
1992. Age, growth, and reproduction of gag, *Mycteroperca microlepis* (Pisces: Serranidae), in the eastern Gulf of Mexico. Bull. Mar. Sci. 51:337–352

Hubble, M.  
2003. The ecological significance of body size in tropical wrasses (Pisces : Labridae). Ph.D diss. James Cook University, Australia.

Killam, K. A., and G. R., Parsons.  
1989. Age and growth of the blacktip shark, *Carcharhinus limbatus*, near Tampa Bay, Florida. Fish. Bull. 87:845–857

Kowal, K.  
2010. Aspects of the life history of the snowy grouper, *Epinephelus niveatus*, in the Gulf of Mexico. M.S. thesis. University of South Florida.

Kulbicki, M., Morize, E., and L. Wantiez.  
2009. Synopsis of the biology and ecology of *Pomadasys argenteus* (Haemulidae) in New Caledonia. Cybium 33:45–59

Lessa, R., and F. M., Santana.  
1998. Age determination and growth of the smalltail shark, *Carcharhinus porosus*, from northern Brazil. Mar. Freshw. Res. 49:705. <https://doi.org/10.1071/MF98019>.

Liu, K.-M., and C.-T., Chen.  
1999. Demographic analysis of the scalloped hammerhead, *Sphyrna lewini*, in the northwestern Pacific. Fish. Sci. 65:218–223

Liu, K.-M., Chiang, P.-J., and C.-T., Chen.  
1998. Age and growth estimates of the bigeye thresher shark, *Alopias superciliosus*, in northeastern Taiwan waters. Fish. Bull. 96:482–491

Loefer, J. K., and G. R., Sedberry.  
2003. Life history of the Atlantic sharpnose shark (*Rhizoprionodon terraenovae*) (Richardson, 1836) off the southeastern United States. S. C. State Doc. Depos.

Lombardi-Carlson, L. A., Fitzhugh, G. R., and J. J., Mikulas.  
2002. Red grouper (*Epinephelus morio*) age-length structure and description of growth from the eastern Gulf of Mexico: 1992-2001. NMFS Southeast. Fish Sci. Cent. Panama City FL Contrib. Ser. 2006:1–37

Lombardi-Carlson, Linda A., Cortes, E., Parsons, G. R., and C. A., Manire. 2003. Latitudinal variation in life-history traits of bonnethead sharks, *Sphyrna tiburo* (Carcharhiniformes: Sphyrnidae) from the eastern Gulf of Mexico. Mar. Freshw. Res. 54:875–883. <https://doi.org/10.1071/MF03023>.

Loubens, G. 1980. Biologie de quelques espèces de poissons du lagon néo-calédonien. II. Sexualité et reproduction. In Cah. L'Indo-Pac. Vol 2. p. 41–72.

Mackie, M. 2000. Reproductive biology of the halfmoon grouper, *Epinephelus rivulatus*, at Ningaloo Reef, Western Australia. Environ. Biol. Fishes 57:363–376

Mahé, K., Rabhi, K., Bellamy, E., Elleboode, R., Aumond, Y., Huet, J., Cresson, P., and D., Roos. 2016. Growth of the oblique-banded grouper (*Epinephelus radiatus*) on the coasts of Reunion Island (SW Indian Ocean). Cybium 40:61–65

Mahmoud, H. H. 2009. Gonadal maturation and histological observations of *Epinephelus areolatus* and *Lethrinus nebulosus* in Halaieb/Shalatien Area “Red Sea”, Egypt. Glob. Vet. 3:414–423

Manickchand-Heileman, S. C., and D. A. T., Phillip. 2000. Age and growth of the yellowedge grouper, *Epinephelus flavolimbatus*, and the yellowmouth grouper, *Mycteroperca interstitialis*, off Trinidad and Tobago. Fish. Bull. 98:290–298

Manooch, C. S. 1987. Age and Growth of Snappers and Groupers. In Trop. Snappers Groupers Biol. Fish. Manag. (Polovina, J.J., and S. Ralston, eds). p. 329–374. Westview Press, Boulder and London.

Mapleston, A., Currey, L. M., Williams, A. J., Pears, R., Simpfendorfer, C. A., Penny, A. L., Tobin, A., and D., Welch. 2009. Comparative biology of key inter-reefal serranid species on the Great Barrier Reef. Proj. Milest. Rep. Mar. Trop. Sci. Res. Facil. Reef Rainfor. Res. Cent. Ltd. Cairns 55 Pp

Mathews, C. P., and M., Samuel.  
1991. Growth, mortality and length-weight parameters for some Kuwaiti fish and shrimp.  
Fishbyte 30–33

Mattos, S. M., Broadhurst, M., Hazin, F. H., and D. M., Jonnes.  
2001. Reproductive biology of the Caribbean sharpnose shark, *Rhizoprionodon porosus*, from northern Brazil. Mar. Freshw. Res. 52:745–752

McBride, R. S., Thurman, P. E., and L. H., Bullock.  
2008. Regional variations of hogfish (*Lachnolaimus maximus*) life history: Consequences for spawning biomass and egg production models. J. Northwest Atl. Fish. Sci. 41:1–12.  
<https://doi.org/10.2960/J.v41.m609>.

McBride, R. S., and A. K., Richardson.  
2007. Evidence of size-selective fishing mortality from an age and growth study of hogfish (Labridae: *Lachnolaimus maximus*), a hermaphroditic reef fish. Bull. Mar. Sci. 80:401–417

McBride, R. S., Sulak, K. J., Thurman, P. E., and A. K., Richardson.  
2009. Age, growth, mortality, and reproduction of roughtongue bass, *Pronotogrammus martinicensis* (Serranidae), in the northeastern Gulf of Mexico. Gulf Mex. Sci. 27:30–38

McGovern, J. C., Wyanski, D. M., Pashuk, O., Manooch, C. S., and G. R., Sedberry.  
1998. Changes in the sex ratio and size at maturity of gag, *Mycteroperca microlepis*, from the Atlantic coast of the southeastern United States. Fish. Bull. 96:797–807

Mees, C. C.  
1992. Seychelles demersal fishery: An analysis of data relating to four key demersal species (Technical Report). Mahe, Seychelles: Seychelles Fishing Authority.

Mehanna, S. F., Al-Marzouqi, A., and B., El-Siabi.  
2013. Stock characteristics and population dynamics of the spiny cheek grouper *Epinephelus diacanthus* (Valenciennes, 1828) from the Arabian Sea, Oman. Turk. J. Fish. Aquat. Sci. 13

Morales-Nin, B.  
1994. Growth of demersal fish species of the Mexican Pacific Ocean. Mar. Biol. 121:211–217

Nardi, K., Newman, S. J., Moran, M. J., and G. P., Jones.  
2006. Vital demographic statistics and management of the baldchin groper (*Choerodon rubescens*) from the Houtman Abrolhos Islands. Mar. Freshw. Res. 57:485.  
<https://doi.org/10.1071/MF05187>.

Neer, J. A., Thompson, B. A., and J. K., Carlson.  
2005. Age and growth of *Carcharhinus leucas* in the northern Gulf of Mexico: incorporating variability in size at birth. J. Fish Biol. 67:370–383

Nichols, R. S., and E. E., DeMartini.  
2008. Preliminary estimates of age and growth for the endemic Hawai’ian grouper (Hapu ‘upu ‘u, *Epinephelus quernus*, F. Serranidae). Pacific Islands Fisheries Science Center, National Marine Fisheries Service, NOAA Honolulu, Pacific Islands Fisheries Science Center Adm. Rep. H-08-06

Ohta, I., and A., Ebisawa.  
2016. Age-based demography and sexual pattern of the white-streaked grouper, *Epinephelus ongus* in Okinawa. Environ. Biol. Fishes 99:741–751. <https://doi.org/10.1007/s10641-016-0517-6>.

Parsons, G. R.  
1985. Growth and age estimation of the Atlantic sharpnose shark, *Rhizoprionodon terraenovae*: A comparison of techniques. Copeia 1985:80. <https://doi.org/10.2307/1444793>.

Parsons, G. R.  
1993. Geographic variation in reproduction between two populations of the bonnethead shark, *Sphyrna tiburo*. In Reprod. Dev. Sharks Skates Rays Ratfishes. p. 25–35. Springer.

Pastor, O. T., Stefánsson, G., Taylor, L., Hjörleifsson, E., and B. Æ., Steinarsson  
2002. Life history and stock assessment of the African hind (*Cephalopholis taeniops*) (Valenciennes, 1828) in São Vicente-São Nicolau insular shelf of the Cape Verde archipelago. Fish. Train. Programme U. N. Univ.

Pears, R. J., Choat, J. H., Mapstone, B. D., and G. A., Begg.  
2006. Demography of a large grouper, *Epinephelus fuscoguttatus*, from Australia’s Great Barrier Reef: implications for fishery management. Mar. Ecol. Prog. Ser. 307:259–272

Piercy, A. N., Carlson, J. K., and M. S., Passerotti.  
2010. Age and growth of the great hammerhead shark, *Sphyrna mokarran*, in the north-western Atlantic Ocean and Gulf of Mexico. Mar. Freshw. Res. 61:992.  
<https://doi.org/10.1071/MF09227>.

Piercy, A. N., Carlson, J. K., Sulikowski, J. A., and G. H., Burgess.  
2007. Age and growth of the scalloped hammerhead shark, *Sphyrna lewini*, in the north-west Atlantic Ocean and Gulf of Mexico. Mar. Freshw. Res. 58:34.  
<https://doi.org/10.1071/MF05195>.

Pitt, J. M., Trott, T. M., and B. E., Luckhurst.  
2010. Bluestriped grunt (*Haemulon sciurus*) in Bermuda: Age, growth, and reproduction studies. In Proc. Gulf Caribb. Fish. Inst. p. 241–245. Gulf and Caribbean Fisheries Institute, c/o Harbor Branch Oceanographic Institution, Inc. Fort Pierce FL 34946 United States., Vol. 62

Pothin, K., Letourneur, Y., and R., Lecomte-Finiger.  
2004. Age, growth and mortality of the tropical grouper *Epinephelus merra* (Pisces, Serranidae) on Reunion Island, SW Indian Ocean. Vie Milieu 54:193–202

Potts, J. C., and C.S., Manooch.  
1999. Observations on the age and growth of graysby and coney from the Southeastern United States. Trans. Am. Fish. Soc. 128:751–757.

Rhodes, K., Taylor, B., and J., McIlwain.  
2011. Detailed demographic analysis of an *Epinephelus polyphekadion* spawning aggregation and fishery. Mar. Ecol. Prog. Ser. 421:183–198. <https://doi.org/10.3354/meps08904>.

Robbins, W. D.  
2006. Abundance, demography and population structure of the grey reef shark (*Carcharhinus amblyrhynchos*) and the white tip reef shark (*Triaenodon obesus*) (Fam. Charcharhinidae). Ph.D diss. James Cook University, Australia.

Romine, J. G., Grubbs, R. D., and J. A., Musick.  
2006. Age and growth of the sandbar shark, *Carcharhinus plumbeus*, in Hawaiian waters through vertebral analysis. Environ. Biol. Fishes 77:229–239. <https://doi.org/10.1007/s10641-006-9113-5>.

Sadovy, Y., and P. L., Colin.  
1995. Sexual development and sexuality in the Nassau grouper. *J. Fish Biol.* 46:961–976

Sadovy, Y. J., Punt, A. E., Cheung, W., Vasconcellos, M., Suharti, S., and Mapstone, B. D.  
2007. Stock assessment approach for the Napoleon fish, *Cheilinus undulatus*, in Indonesia: A tool for quota-setting for data-poor fisheries under CITES Appendix II Non-Detriment Finding requirements. Food and Agriculture Organization of the United Nations.

Sadovy, Y. J., Figuerola, M., and A., Roman.  
1992. Age and growth of red hind *Epinephelus guttatus* in Puerto Rico and St. Thomas. *Fish. Bull.* 90:516–528

Semba, Y. J., Nakano, H., and I. Aoki.  
2009. Age and growth analysis of the shortfin mako, *Isurus oxyrinchus*, in the western and central North Pacific Ocean. *Environ. Biol. Fishes* 84:377–391.  
<https://doi.org/10.1007/s10641-009-9447-x>.

Simpfendorfer, C. A.  
1992. Reproductive strategy of the Australian sharpnose shark, *Rhizoprionodon taylori* (Elasmobranchii: Carcharhinidae), from Cleveland Bay, northern Queensland. *Mar. Freshw. Res.* 43:67–75

Simpfendorfer, C. A.  
1993. Age and growth of the Australian sharpnose shark, *Rhizoprionodon taylori*, from north Queensland, Australia. *Environ. Biol. Fishes* 36:233–241

Simpfendorfer, C. A., McAuley, R. B., Chidlow, J., and P., Unsworth.  
2002. Validated age and growth of the dusky shark, *Carcharhinus obscurus*, from Western Australian waters. *Mar. Freshw. Res.* 53:567–573

Sujatha, K., and K. V. L., Shrikanya.  
2013. Reproductive biology of striped grouper *Epinephelus latifasciatus* (Temminck and Schelegel, 1842), off Visakhapatnam, middle east coast of India. *Indian J. Geo-Mar. Sci.* 42:183–190

Tariche, O., Pajuelo, J. G., Lorenzo, J. M., Luque, A., and J. A., Gonzalez. 2015. Age estimation and growth pattern of the grouper *Cephalopholis taeniops* (Epinephelidae) off the Cape Verde Archipelago, north-west Africa. *J. Mar. Biol. Assoc. U. K.* 95:599–609. <https://doi.org/10.1017/S0025315414001441>.

Torres Jr., F., and Pauly, D. 1991. Tabular data on marine fishes from Southern Africa, Part II: Growth parameters. *Fishbyte* 9:37–38

Trott, T. M. 2006. Preliminary analysis of age, growth, and reproduction of coney (*Cephalopholis fulva*) at Bermuda. *57th Gulf and Caribbean Fisheries Institute*. 385–400.

Usseglio, P., Friedlander, A. M., DeMartini, E. E., Schuhbauer, A., Schemmel, E., and P., Salinas de Léon. 2015. Improved estimates of age, growth and reproduction for the regionally endemic Galapagos sailfin grouper *Mycteroperca olfax* (Jenyns, 1840). *PeerJ* 3:e1270. <https://doi.org/10.7717/peerj.1270>.

Valle, S. V., Garcia-Arteaga, J. P., and R., Claro. 1997. Growth parameters of marine fishes in Cuban waters. *Naga*. 34–37

Wakefield, C. B., Newman, S. J., Marriott, R. J., Boddington, D. K., and D. V., Fairclough. 2013. Contrasting life history characteristics of the eightbar grouper *Hyporthodus octofasciatus* (Pisces: Epinephelidae) over a large latitudinal range reveals spawning omission at higher latitudes. *ICES J. Mar. Sci.* 70:485–497. <https://doi.org/10.1093/icesjms/fst020>.

Wenjia, H., Guanqiong, Y., Zhenbin, L., Jianguo, D., Mingru, C., Loke Ming, C., and Y., Shengyun. 2015. Study on fish life history traits and variation in the Taiwan Strait and its adjacent waters. *Acta Oceanol. Sin.* 34:45–54. <https://doi.org/10.1007/s13131-015-0625-8>.

White, W. T., Hall, N., and I., Potter. 2002. Size and age compositions and reproductive biology of the nervous shark *Carcharhinus cautus* in a large subtropical embayment, including an analysis of growth during pre- and postnatal life. *Mar. Biol.* 141:1153–1164. <https://doi.org/10.1007/s00227-002-0914-6>.

Williams, A., Davies, C., Mapstone, B., Currey, L., Welch, D., Begg, G., Ballagh, A., Choat, J., Murchie, C., and C., Simpfendorfer.

2009. Age-based demography of humpback grouper *Cromileptes altivelis*: implications for fisheries management and conservation. *Endanger. Species Res.* 9:67–79.  
<https://doi.org/10.3354/esr00237>.

Wyanski, D. M., White, D. B., and C. A. Barans.

2000. Growth, population age structure, and aspects of the reproductive biology of snowy grouper, *Epinephelus niveatus*, off North Carolina and South Carolina. S. C. State Doc. Depos.